

A FIRST ESTIMATE OF THE BARYONIC MASS FUNCTION OF GALAXIES

ERIC F. BELL

Max-Planck-Institut für Astronomie, Königstuhl 17, D-69117 Heidelberg, Germany; bell@mpia.de

AND

DANIEL H. MCINTOSH, NEAL KATZ, AND MARTIN D. WEINBERG

Department of Astronomy, University of Massachusetts at Amherst, 640 Lederle Graduate Research Center, Amherst, MA 01003;

dmac@hamerkop.astro.umass.edu, nsk@kaka.astro.umass.edu, weinberg@astro.umass.edu

Received 2002 December 13; accepted 2003 January 30; published 2003 February 7

ABSTRACT

We estimate the baryonic (stellar+cold gas) mass function of galaxies in the local universe by assigning a complete sample of Two Micron All Sky Survey and Sloan Digital Sky Survey galaxies a gas fraction based on a statistical sample of the entire population, under the assumption of a universally applicable stellar initial mass function. The baryonic mass function is similar to the stellar mass function at the high-mass end and has a reasonably steep faint-end slope owing to the typically high cold gas fractions and low stellar mass-to-light ratios characteristic of low-mass galaxies. The Schechter function fit parameters are $\phi^* h^{-3} = 0.0108(6) \text{ Mpc}^{-3} \log M^{-1}$, $M^* h^2 = 5.3(3) \times 10^{10} M_{\odot}$, and $\alpha = -1.21(5)$, with formal error estimates given in parentheses. We show that the H I and H₂ mass functions derived using this indirect route are in agreement with direct estimates, validating our indirect method. Integrating under the baryonic mass function and incorporating all sources of uncertainty, we find that the baryonic (stellar+cold gas) mass density implied by this estimate is $\Omega_{\text{cold baryon}} h = 2.4_{-1.4}^{+0.7} \times 10^{-3}$, or $8_{-3}^{+4}\%$ of the big bang nucleosynthesis expectation.

Subject headings: galaxies: general — galaxies: ISM — galaxies: luminosity function, mass function — galaxies: stellar content

1. INTRODUCTION

The distribution of the mass in collapsed baryons (cold gas and stars) in galaxies is a fundamental prediction of galaxy formation models. Unfortunately, to date there is no robust estimate of the baryonic mass function (MF) of galaxies, leaving modelers with the nontrivial task of predicting stellar masses or, even worse, galaxy luminosities. Discrepancy between the model and data may indicate a problem with the predicted distribution of galaxy baryonic masses or could represent poorly constrained star formation (SF), stellar population, or dust prescriptions. In this Letter, we present a first estimate of the baryonic MF of galaxies by assigning galaxies' gas fractions statistically (based on an independent sample), under the assumption of a universally applicable stellar initial mass function (IMF).¹

The time is ripe to attempt this for the first time. With the advent of large, relatively complete surveys, the luminosity function (LF) is now well constrained in the optical and near-infrared (NIR; Gardner et al. 1997; Cole et al. 2001; Kochanek et al. 2001; Norberg et al. 2002; Liske et al. 2002; Blanton et al. 2002; Bell et al. 2003). Furthermore, under the assumption of a universally applicable stellar IMF, the distribution of stellar masses is reasonably well constrained, with an overall normalization uncertainty caused by our relatively poor knowledge of the faint-end slope of the IMF (Cole et al. 2001; Bell et al. 2003). Crucially, there are also relatively large samples of galaxies with *K*-band data and gas masses, allowing a reasonably accurate characterization of the gas mass of galaxies as a function of their physical parameters (Bell & de Jong 2000).

¹ A time-varying IMF, as speculated on by Hernandez & Ferrara (2001) or Ferguson, Dickinson, & Papovich (2002), would invalidate this assumption.

2. THE DATA AND METHODOLOGY

Because of the lack of a large galaxy survey with both gas mass and *K*-band data, we take a sampling approach, analogous to that used by Loveday (2000) to estimate the *K*-band LF from a *B*-band limited survey. Essentially, we estimate a stellar MF (§ 2.1) and then add representative gas masses to each galaxy (§ 2.2), allowing us to estimate the distribution of galaxy baryonic masses (§ 3). We assume $\Omega_m = 0.3$, $\Omega_{\Lambda} = 0.7$, and $H_0 = 100 h \text{ km s}^{-1} \text{ Mpc}^{-1}$.

2.1. Estimating the Stellar Mass Function

We construct the baryonic MF using a combined sample of galaxies from the Two Micron All Sky Survey (2MASS; Skrutskie et al. 1997) and the Sloan Digital Sky Survey (SDSS; York et al. 2000). We use the SDSS early data release (EDR; Stoughton et al. 2002) to provide an 84% redshift complete $r \leq 17.5$ sample of galaxies with accurate *ugriz* fluxes over 414 deg², which is ~10% less than the whole EDR imaging area because some spectroscopic plates were not attempted (Stoughton et al. 2002). The 84% redshift completeness within this area is our own direct estimate based on the fraction of galaxies fulfilling the Strauss et al. (2002) criteria that have spectra, in agreement with the EDR analysis of Nakamura et al. (2003). To account for light missed by the Petrosian magnitude estimator (Strauss et al. 2002; Blanton et al. 2002), we add 15% to the fluxes of galaxies morphologically classified as early type using the SDSS *r*-band concentration parameter following Strateva et al. (2001). This correction produces only a $\lesssim 5\%$ effect on the LFs and MFs (Blanton et al. 2002; Bell et al. 2003). We also correct for an ~8% overdensity of galaxies in the EDR, as estimated by comparing the number of $10 \leq K \leq 13.5$ galaxies in the EDR spectroscopic area with that from the sky with $|b| \geq 30^\circ$.

We use the now complete 2MASS extended source and point-source catalogs to augment the SDSS *ugriz* fluxes with *K*-band




FIG. 1.— K -band magnitude and half-light radius for the 6573 late-type SDSS+2MASS galaxies in our sample (*contours*) and for the 156 galaxy comparison sample of Bell & de Jong (2000; *filled circles*).

fluxes and for extended sources K -band half-light radii. We correct 2MASS K -band fluxes to total following a comparison with deeper K -band data from Loveday (2000); for extended sources, this amounts to a 0.1 mag correction (Bell et al. 2003). We do not use 2MASS J - or H -band data here because we cannot correct the magnitudes similarly. The optical and NIR magnitude zero points are accurate to ~ 0.05 and ~ 0.02 mag, respectively, and the random errors are 0.05 mag (optical) and 0.2 mag (NIR).

To estimate k -corrections, evolution corrections, and the present-day stellar mass-to-light (M/L) ratios, we fit the $ugrizK$ observed fluxes² with model stellar populations. These populations have a range of metallicities and SF histories at a given redshift. We use the PÉGASE stellar population synthesis model (see Fioc & Rocca-Volmerange 1997 for a description of an earlier version of their model) with a “diet” Salpeter IMF (following Bell & de Jong 2001) that has the same colors and luminosity as a normal Salpeter IMF but with only 70% of the mass (owing to a smaller number of low-mass stars). Corrections derived using this technique are consistent with those used by Blanton et al. (2002). The stellar M/L ratios we derive are within 10% of those from the spectral modeling technique of Kauffmann et al. (2003a), accounting for differences in IMF; the random and systematic uncertainties from dust and bursts of SF dominate, however, and are $\lesssim 25\%$ (Bell & de Jong 2001). This IMF is “maximum disk,” inasmuch as IMFs richer in faint low-mass stars overpredict the rotation velocity of Ursa Major Cluster galaxies with K -band photometry and well-resolved H I rotation curves. This prescription thus gives the maximum possible stellar M/L ratio. Naturally, a different choice of IMF allows lower M/L ratios. For example, the popular Kennicutt or Kroupa IMFs have $\sim 30\%$ lower M/L ratios than this IMF and are thus “sub-maximal” (see Bell & de Jong 2001 for more discussion of this point).

We calculate LFs and stellar MFs using the V/V_{\max} formalism (Felten 1977), taking into account foreground Galactic extinction, k -corrections, and evolution corrections. In Bell et al. (2003), we match precisely published LFs; in particular, we

² Not all galaxies have $ugrizK$ fluxes. We have checked that missing passbands do not significantly bias the estimated k -corrections, evolution corrections, or stellar M/L ratios (but do, of course, increase the random error somewhat).

reproduce the g -band and K -band LF and luminosity densities to within $\lesssim 10\%$ (Blanton et al. 2002; Cole et al. 2001; Kochanek et al. 2001). Furthermore, this method produces accurate stellar MFs that match the estimate of Cole et al. (2001) to $\sim 10\%$ in total stellar mass density (accounting for IMF differences), *but can do so using LFs limited by optical or NIR magnitude limits* (Bell et al. 2003). For this Letter, we choose 11,848 galaxies with $13 \leq r \leq 17.5$ and $g \leq 17.74$, which ensures that we have accurate $g-r$ color estimates providing a stellar M/L ratio accuracy of better than 25%, while avoiding potential biases against low surface brightness galaxies in 2MASS (Bell et al. 2003). The stellar MF estimated using this technique is shown in Figure 2, along with the stellar MF of Cole et al. (2001) for comparison. A much more extensive description of the LF and stellar MF construction is given by Bell et al. (2003).

2.2. Estimating Gas Masses

Because there are no samples of galaxies with good number statistics, deep optical/NIR data, and gas masses, we estimate the gas masses of SDSS+2MASS galaxies indirectly. We use galaxies from Bell & de Jong (2000) with K -band luminosities, half-light radii, and gas masses to statistically assign a gas mass to every SDSS+2MASS galaxy, appropriate to its K -band luminosity and half-light radius.

Figure 1 shows the K -band half-light radii and luminosities for the late-type subsample of the SDSS+2MASS galaxies (*contours*) and for the comparison sample of 156 galaxies with gas masses (*filled circles*) taken from Bell & de Jong (2000). We estimate gas masses by multiplying the H I gas mass by 1.33 to account for helium and by the morphological type-dependent H_2 -to-H I ratio presented by Young & Knezek (1989). For the 21% of the galaxies without K -band luminosities and sizes, we adopt the r -band half-light radii and estimate the K -band luminosity by dividing the g -band-derived stellar mass by the model K -band stellar M/L ratio. Our assumptions are accurate to better than 40% in both cases and make no difference to our results.

Each SDSS+2MASS galaxy is assigned a gas mass from a galaxy in the Bell & de Jong (2000) comparison sample with similar half-light radius and K -band luminosity. For galaxies morphologically classified as late type, we assign gas masses using a randomly chosen comparison galaxy that is within a factor of 2 in size and within 1 mag in K -band luminosity. Galaxies are morphologically classified using the r -band concentration parameter following Strateva et al. (2001). We scale the gas mass by the difference in luminosity to conserve the galaxy gas fraction. As a consistency check, we also assign gas masses by choosing the nearest galaxy in half-light radius–luminosity space and by assigning gas to galaxies of all morphological types (see Fig. 2). These changes make no appreciable difference to our results.

3. THE BARYONIC GALAXY MASS FUNCTION

Our baryonic galaxy MF is shown as the solid line with open circles in the left panel of Figure 2. Shown for comparison is the stellar MF from Cole et al. (2001) and the g -selected stellar MF described above and in Bell et al. (2003). The baryonic galaxy MF follows the stellar MF at high masses, shows a modest offset at the “knee” of the MF, and shows a reasonably steep faint-end slope. This behavior is expected, as low-mass field galaxies tend to have high cold gas fractions and more ongoing SF (Bell & de Jong 2000; Kauffmann et al. 2003b),




FIG. 2.—Baryonic MF of galaxies. *Left panel:* Stellar MF of Cole et al. (2001; *naked error bars*) corrected to our maximum disk IMF, the stellar MF derived using our SDSS *g*-band-selected sample (*solid gray line*), and the baryonic MF of galaxies, assuming the maximum disk IMF (*solid line with open circles and error bars*, with a Schechter fit as the thin solid line). *Middle panel:* Different versions of the baryonic MF, illustrating the different sources of uncertainty. The solid line with open circles again is the baryonic MF defined using the default gas mass estimation technique, the dashed line shows the effect of choosing the nearest galaxy on the half-light radius–luminosity plane for estimating the gas mass, the dotted line shows the effect of allowing SDSS+2MASS early-type galaxies to also have gas, and the two solid lines are the effect of choosing a Kennicutt or Kroupa IMF (*upper line*) or an *M/L* ratio consistent with a Bottema disk (*lowest line*). *Right panel:* We test the gas mass estimation method by comparing the H I and H₂ MFs predicted using this method with observations. The solid line with open circles and error bars denotes the match of SDSS+2MASS spiral galaxies with randomly selected nearby galaxies, and the dotted and dashed lines are as in the middle panel. The solid black curve is the blind H I MF of Rosenberg & Schneider (2002). The H₂ galaxy MF is shown as the gray line with open circles and error bars, and the Schechter fit to the first observational estimate of the H₂ MF (from Keres et al. 2003) is shown as the solid gray line.

which steepens the baryonic MF compared to optical/NIR LFs that typically have $\alpha \sim -1.0$. The Schechter function fit parameters are $\phi^* h^{-3} = 0.0108(6) \text{ Mpc}^{-3} \log M^{-1}$, $M^* h^2 = 5.3(3) \times 10^{10} M_{\odot}$, and $\alpha = -1.21(5)$, where the formal error estimates are in parentheses.

In the middle panel of Figure 2, we show the effects of the systematic uncertainties. In particular, the dashed, dotted, and dot-dashed lines show the effects of changing the gas mass assignment (see the figure legend for more details); these uncertainties have only small effects. The two bare solid lines show the effect of assuming different stellar IMFs. As stated earlier, we adopt as our default an IMF that has the largest stellar *M/L* ratio permitted, without overpredicting the rotation velocities of spiral galaxies in the Ursa Major Cluster (*solid line with open circles*; Bell & de Jong 2001). Yet, the stellar *M/L* ratio may be lower than this maximal value; thus, we show two cases. First, we plot (*upper solid line*) the increasingly popular Kroupa, Tout, & Gilmore (1993) and Kennicutt (1983) IMFs, which both have *M/L* ratios of $\sim 70\%$ of maximal IMF ($\sim 50\%$ Salpeter). The other case is an IMF that has *M/L* ratios of only 40% of our maximal IMF (*lowest solid line*), which corresponds roughly to the 63% disk velocity contribution to the rotation curve as argued by Bottema (1997). It is clear that, assuming a universal IMF, the factor of 2 uncertainty in the stellar *M/L* ratio dominates the error budget in terms of total cold baryonic mass in the local universe. Given the stellar IMF, the systematic uncertainties are $\lesssim 25\%$. We estimate these uncertainties using different passbands to estimate stellar mass, using different gas mass assignment methods, and accounting for the effects of dust and bursts of SF on the *M/L* ratios (Bell & de Jong 2001; Bell et al. 2003). The Schechter function fit parameters for the Kennicutt/Kroupa case and the Bottema case are: $\phi^* h^{-3} = 0.0116(5)$, $0.0142(8) \text{ Mpc}^{-3} \log M^{-1}$, $M^* h^2 = 3.78(11)$, $2.24(8) \times 10^{10} M_{\odot}$, and $\alpha = -1.22(3)$, $-1.20(3)$.

It is important to make sure that our statistical procedure

assigns gas masses consistent with the true galaxy population by comparing with the H I or H₂ MF of galaxies (Fig. 2, *right panel*). The solid black line with open circles and error bars is our H I MF derived in this way. The dashed and dotted lines show the effects of using the closest galaxy to estimate gas mass and allowing elliptical galaxies to have gas mass, respectively. For comparison, the H I MF of the blind H I Arecibo survey of Rosenberg & Schneider (2002) is plotted as the smooth solid curve. We also show our prediction of the H₂ galaxy MF as the solid gray line with open circles and error bars. For comparison, we plot the Schechter fit to the first observational estimate of the H₂ MF (Keres, Yun, & Young 2003). Given the uncertainties, it is clear that our method for estimating galaxy gas masses reproduces the H I and H₂ galaxy MFs with astonishing precision. Therefore, the sample of galaxies that we use to assign gas masses is indeed reasonably representative and lends considerable credibility to our estimate of the baryonic galaxy MF.

4. DISCUSSION

Even with the factor of 2 uncertainty from the contribution of low-mass stars to the overall stellar *M/L* ratio, we can still draw some conclusions about the local universe. It is clear that the overall efficiency of galaxy formation is very low. First, the faint-end slope of the baryonic MF is ~ -1.2 , which is much shallower than the ~ -2 expected for the halo MF (e.g., White & Frenk 1991). Second, integrating under the MF, we derive $\Omega_{\text{cold baryon}} h = 2.4^{+0.7}_{-1.4} \times 10^{-3}$, including the IMF and 25% systematic stellar *M/L* ratio uncertainties. Our estimate agrees well with the value of $(2.9 \pm 1.5) \times 10^{-3}$ from Fukugita, Hogan, & Peebles (1998) and is preferred owing to our better accounting for stellar *M/L* ratios compared with Fukugita et al. (1998), who use (harder to convert into stellar mass) *B*-band luminosity densities assuming a similar IMF to the

maximum-disk IMF we adopt here. Taking the value of the total baryon density from the big bang nucleosynthesis value of O’Meara et al. (2001), and assuming $h = 0.7 \pm 0.07$, we find $\Omega_{\text{cold baryon}}/\Omega_b \sim 8_{-5}^{+4}\%$, where the error estimates account for the uncertainties in IMF, H_0 , Ω_b , our gas assignment method, and the $\lesssim 25\%$ uncertainties in stellar M/L ratios from dust and bursts of SF. Our value is quite consistent with the low galaxy formation efficiency characteristic of most current models, which have low efficiencies at the low- and high-mass ends because of feedback from supernovae and inefficient gas cooling, respectively (e.g., Cole et al. 2000).

Accounting for the possible gaseous content of elliptical galaxies, for submaximal M/L ratios, and for the effects of dust and bursts of SF on stellar M/L ratios, the universal gas fraction, $f_g = \Omega_{\text{cold gas}}/\Omega_{\text{cold baryons}}$, should lie in the range $0.2 \leq f_g \leq 0.5$. For the “maximal” IMF, we find $f_g \sim 25\%$. Fukugita et al. (1998) and Keres et al. (2003) find values of 15%–20% when our IMF is adopted; their slightly lower determinations stem primarily from a lower estimate of H I mass density. Nevertheless, all the studies agree that $f_g \leq 0.5$; therefore, the dynamically cold baryons (i.e., the gas and stars in disks and spheroids) are primarily in the form of stars, even for low stellar M/L ratios.

It is well known observationally that cluster optical/NIR LFs have steeper faint-end slopes than field LFs (e.g., Trentham & Tully 2002). Furthermore, cluster galaxies tend to have little ongoing SF and little gas, so that most cluster galaxies are star dominated with large stellar M/L ratios (e.g., Kuntschner 2000). Thus, the trend of increasing faint-end slope with increasing cluster mass noted by, e.g., Trentham & Tully (2002) may be more naturally interpreted as a constant baryonic MF, with a suppression of recent SF in massive clusters of galaxies. Obviously, a deeper investigation of this issue is warranted before speculating any further.

5. CONCLUSIONS

Together with the baryonic (stellar+cold gas) luminosity–line width relation (e.g., McGaugh et al. 2000; Bell & de Jong 2001), the baryonic galaxy MF is an ideal test of models of galaxy formation and evolution. In this Letter, we have estimated the baryonic galaxy MF in the local universe for the first time assuming a universally applicable stellar IMF. We assign gas fractions statistically to a large sample of galaxies from 2MASS and SDSS, using a local sample with accurate K -band and gas fraction data. We cross-check this statistical procedure against independent H I and H₂ surveys, finding excellent agreement. The baryonic MF is similar to the stellar MF at the high-mass end (with a slightly higher density normalization) and has a reasonably steep faint-end slope, $\alpha \sim -1.2$, due to the typically high cold gas fractions and low stellar M/L ratios of low-mass galaxies. Integrating under the baryonic MF, we find that the baryonic (stellar+cold gas) mass density implied by this estimate is $\Omega_{\text{cold baryon}} h = 2.4_{-1.4}^{+0.7} \times 10^{-3}$, or $8_{-5}^{+4}\%$ of the big bang nucleosynthesis expectation. This clearly implies a low overall efficiency of galaxy formation.

E. F. B. was supported by the European Research Training Network “Spectroscopic and Imaging Surveys for Cosmology.” D. H. M. and M. D. W. acknowledge support by JPL/NASA through the 2MASS core science projects. This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation. This publication also makes use of the Sloan Digital Sky Survey.³ We thank the referee, Andrea Ferrara, for his helpful suggestions and feedback.

³ See <http://www.sdss.org>.

REFERENCES

- Bell, E. F., & de Jong, R. S. 2000, *MNRAS*, 312, 497
 ———. 2001, *ApJ*, 550, 212
 Bell, E. F., McIntosh, D. H., Katz, N., & Weinberg, M. D. 2003, *ApJ*, submitted
 Blanton, M. R., et al. 2002, *ApJ*, submitted (astro-ph/0210215)
 Bottema, R. 1997, *A&A*, 328, 517
 Cole, S., Lacey, C., Baugh, C. M., & Frenk, C. S. 2000, *MNRAS*, 319, 168
 Cole, S., et al. 2001, *MNRAS*, 326, 255
 Felten, J. E. 1977, *AJ*, 82, 861
 Ferguson, H. C., Dickinson, M., & Papovich, C. 2002, *ApJ*, 569, L65
 Fioc, M., & Rocca-Volmerange, B. 1997, *A&A*, 326, 950
 Fukugita, M., Hogan, C. J., & Peebles, P. J. E. 1998, *ApJ*, 503, 518
 Gardner, J. P., Sharples, R. M., Frenk, C. S., & Carrasco, B. E. 1997, *ApJ*, 480, L99
 Hernandez, X., & Ferrara, A. 2001, *MNRAS*, 324, 484
 Kauffmann, G., et al. 2003a, *MNRAS*, in press (astro-ph/0204055)
 ———. 2003b, *MNRAS*, in press (astro-ph/0205070)
 Kennicutt, R. C., Jr. 1983, *ApJ*, 272, 54
 Keres, D., Yun, M. S., & Young, J. S. 2003, *ApJ*, 582, 659
 Kochanek, C. S., et al. 2001, *ApJ*, 560, 566
 Kroupa, P., Tout, C. A., & Gilmore, G. 1993, *MNRAS*, 262, 545
 Kuntschner, H. 2000, *MNRAS*, 315, 184
 Liske, J., Lemon, D. J., Driver, S. P., Cross, N. J. G., & Couch, W. J. 2002, *MNRAS*, submitted (astro-ph/0207555)
 Loveday, J. 2000, *MNRAS*, 312, 557
 McGaugh, S. S., Schombert, J. M., Bothun, G. D., & de Blok, W. J. G. 2000, *ApJ*, 533, L99
 Nakamura, O., et al. 2003, *AJ*, in press (astro-ph/0212405)
 Norberg, P., et al. 2002, *MNRAS*, 332, 827
 O’Meara, J. M., et al. 2001, *ApJ*, 552, 718
 Rosenberg, J. L., & Schneider, S. E. 2002, *ApJ*, 567, 247
 Skrutskie, M. F., et al. 1997, in *The Impact of Large Scale Near-IR Sky Surveys*, ed. F. Garzon et al. (Dordrecht: Kluwer), 25
 Stoughton, C., et al. 2002, *AJ*, 123, 485
 Strateva, I., et al. 2001, *AJ*, 122, 1861
 Strauss, M. A., et al. 2002, *AJ*, 124, 1810
 Trentham, N., & Tully, R. B. 2002, *MNRAS*, 335, 712
 White, S. D. M., & Frenk, C. S. 1991, *ApJ*, 379, 52
 York, D. G., et al. 2000, *AJ*, 120, 1579
 Young, J. S., & Knezek, P. M. 1989, *ApJ*, 347, L55